

A COUNTER EXAMPLE TO MALLE'S CONJECTURE ON THE ASYMPTOTICS OF DISCRIMINANTS

JÜRGEN KLÜNERS

ABSTRACT. In this note we give a counter example to a conjecture of Malle which predicts the asymptotic behaviour of the counting functions for field extensions with given Galois group and bounded discriminant.

1. INTRODUCTION

Let $G \leq S_n$ be a finite transitive permutation group and k be a number field. We say that a finite extension K/k has Galois group G if the normal closure $\hat{K}$ of K/k has Galois group isomorphic to G and K is the fixed field in $\hat{K}$ under a point stabilizer of G . By abuse of notation we will write $\text{Gal}(K/k) = G$ in this situation. We let

$$Z(k, G; x) := |\{K/k : \text{Gal}(K/k) = G, N_{k/\mathbb{Q}}(d_{K/k}) \leq x\}|$$

be the number of field extensions of k (inside a fixed algebraic closure $\bar{\mathbb{Q}}$) of degree n with Galois group permutation isomorphic to G (as explained above) and norm of the discriminant $d_{K/k}$ bounded above by x . It is well known that the number of extensions of k with bounded norm of the discriminant is finite, hence $Z(k, G; x)$ is finite for all G , k and $x \geq 1$.

Gunter Malle [7, 8] has given a precise conjecture about the asymptotic behaviour of the function $Z(k, G; x)$ for $x \rightarrow \infty$. In order to state it, we need to introduce some group theoretic invariants of permutation groups.

Definition 1.1. Let $1 \neq G \leq S_n$ be a transitive subgroup acting on $\Omega = \{1, \dots, n\}$.

- (1) For $g \in G$ we define $\text{ind}(g) := n - \text{the number of orbits of } g \text{ on } \Omega$.
- (2) $\text{ind}(G) := \min\{\text{ind}(g) : 1 \neq g \in G\}$.
- (3) $a(G) := \text{ind}(G)^{-1}$.

Since all elements in a conjugacy class C of G have the same index we can define $\text{ind}(C)$ in a canonical way. The absolute Galois group of k acts on the set of conjugacy classes of G via the action on the columns of the complex character table of G . The orbits under this action are called k -conjugacy classes.

Definition 1.2. For a number field k and a transitive subgroup $1 \neq G \leq S_n$ we define:

$$b(k, G) := |\{C : C \text{ } k\text{-conjugacy class of minimal index } \text{ind}(G)\}|.$$

Now we can state the conjecture of Malle[8], where $f(x) \sim g(x)$ is a notation for $\lim_{x \rightarrow \infty} f(x)/g(x) = 1$.

Conjecture 1.3. (Malle) For all number fields k and all transitive permutation groups $1 \neq G$ there exists a constant $c(k, G) > 0$ such that

$$Z(k, G; x) \sim c(k, G) x^{a(G)} \log(x)^{b(k, G)-1},$$

where $a(G)$ and $b(k, G)$ are given as above.

This conjecture is proved for abelian groups [9]. For all number fields k and all nilpotent groups G it is shown in [6] that

$$\limsup_{x \rightarrow \infty} \frac{\log Z(k, G; x)}{\log x} \leq a(G).$$

If we furthermore assume that G is in its regular representation, i.e., we count normal nilpotent number fields, we get:

$$\lim_{x \rightarrow \infty} \frac{\log Z(k, G; x)}{\log x} = a(G).$$

For more results see also the survey articles [1, 2].

2. THE COUNTER EXAMPLE

We present a counter example for $k = \mathbb{Q}$ and the wreath product $G := C_3 \wr C_2 = C_3^2 \rtimes C_2 \leq S_6$ of order 18, where C_n is the cyclic group of order n . In the following we count all field towers $L/K/\mathbb{Q}$ such that $\text{Gal}(L/K) = C_3$ and $\text{Gal}(K/\mathbb{Q}) = C_2$. Therefore the Galois group of $L/\mathbb{Q}$ is one of the groups $C_6, S_3(6), G \leq S_6$, where $S_3(6)$ denotes the group S_3 in its degree 6 representation. Since C_6 is abelian we get from [9] that

$$Z(\mathbb{Q}, C_6; x) \sim c(C_6)x^{1/3}.$$

From the Davenport-Heilbronn theorem [3] we know that

$$Z(\mathbb{Q}, S_3(3); x) \sim c(S_3(3))x.$$

Using the fact that the discriminant of the splitting field of an S_3 -extension is at least the square of the discriminant of the S_3 -extension, we easily get that

$$Z(\mathbb{Q}, S_3(6); x) = O(x^{1/2}).$$

Since extensions with Galois group $S_3(6)$ are normal we can use a result in [4, Prop. 2.8] which states that $Z(\mathbb{Q}, S_3(6); x) = O_\epsilon(x^{3/8+\epsilon})$ for all $\epsilon > 0$. With a more careful analysis we are able to prove that there are constants $c_1(S_3), c_2(S_3) > 0$ such that

$$c_1(S_3)x^{1/3} \leq Z(\mathbb{Q}, S_3(6); x) \leq c_2(S_3)x^{1/3} \quad \text{for } x \text{ large enough.}$$

We remark that the results for $S_3(6)$ and C_6 are as conjectured since we get $a(C_6) = a(S_3(6)) = 1/3$ and $b(\mathbb{Q}, C_6) = b(\mathbb{Q}, S_3(6)) = 1$.

Now we define the counting function corresponding to field towers $L/K/\mathbb{Q}$ as above:

$$\tilde{Z}(\mathbb{Q}, C_3 \wr C_2; x) := |\{L/\mathbb{Q} \mid \exists K : \text{Gal}(L/K) = C_3, [K : \mathbb{Q}] = 2, |d_L| \leq x\}|.$$

We have two conjugacy classes of elements of order 3 in G which have three fixed points. Considered as $\mathbb{Q}$ -conjugacy classes we get only one orbit. In number fields k containing a primitive third root of unity ζ_3 we get two k -conjugacy classes of this type. Therefore $a(G) = 1/2$ and $b(\mathbb{Q}, G) = 1$. Since the counting functions for $S_3(6)$ and C_6 have lower asymptotics we would expect that

$$\tilde{Z}(\mathbb{Q}, C_3 \wr C_2; x) \sim Z(\mathbb{Q}, C_3 \wr C_2; x) \sim c(G)x^{1/2}.$$

Certainly we get a lower estimate for $\tilde{Z}(\mathbb{Q}, C_3 \wr C_2; x)$ if we only count the number fields which contain a fixed quadratic subfield K . We choose $K = \mathbb{Q}(\zeta_3)$ and using $d_L = d_K^3 N(d_{L/K})$ we get for x large enough:

$$\tilde{Z}(\mathbb{Q}, C_3 \wr C_2; x) \geq Z(K, C_3; x/27) \sim c(K, C_3) x^{1/2} \log(x).$$

For the latter we used the fact that $b(K, C_3) = 2$ and that the conjecture is true for abelian groups [9]. This already gives a contradiction to Conjecture 1.3. Since the asymptotics of the counting functions of C_6 and $S_3(6)$ is $O(x^{1/3})$ we get the contradiction for our group G .

Now we introduce a counting function avoiding $\mathbb{Q}(\zeta_3)$.

$$\hat{Z}(\mathbb{Q}, C_3 \wr C_2; x) := |\{L/\mathbb{Q} \mid \exists K \neq \mathbb{Q}(\zeta_3) : \text{Gal}(L/K) = C_3, [K : \mathbb{Q}] = 2, |d_L| \leq x\}|.$$

Using the averaging results for the 3-ranks of the class group of quadratic fields [3] we can prove that

$$\hat{Z}(\mathbb{Q}, C_3 \wr C_2; x) \sim c(C_3 \wr C_2) x^{1/2}$$

as predicted by Malle's conjecture. This means that the cyclotomic intermediate extension is the reason for the failure of the conjecture.

We remark that all groups of type $C_\ell \wr C_m$, where $\ell > 2$ is prime and $\gcd(m, \ell - 1) > 1$ give counter examples. E.g. let us consider $m = 2$. Then there is a unique quadratic subfield $K = \mathbb{Q}(\sqrt{\pm \ell})$ of $\mathbb{Q}(\zeta_\ell)$. Since $b(K, C_\ell) = 2$ and $b(\mathbb{Q}, C_\ell \wr C_2) = 1$ we can derive the same contradiction as above. We remark that for $\ell > 3$ we cannot prove good upper bounds for those groups since we do not know good estimates for the ℓ -rank of the class group of quadratic fields in these cases.

How to fix the conjecture?

One possibility would be to forbid intermediate extensions which are contained in cyclotomic extensions $\mathbb{Q}(\zeta_\ell)$, where ℓ must be chosen from a set containing all orders of elements of G which have minimal index. This is more natural in the global function field setting. Here we restrict to extensions $K/\mathbb{F}_q(t)$ such that the normal closure of K contains no constant field extension. Restricting to those extensions and assuming some heuristic about the number of points of irreducible varieties over $\mathbb{F}_q$, Ellenberg and Venkatesh [5] are able to deduce the correct upper and lower bounds in Malle's conjecture including the correct logarithmic power. We remark that the same type of counter examples apply, if we allow constant field extensions. E.g. choosing $q \equiv 2 \pmod{3}$ and $G = C_3 \wr C_2$ works as a counter example, when we choose $\mathbb{F}_{q^2}/\mathbb{F}_q$ as the quadratic extension.

In order to fix the conjecture we might think that we have to look at maximal abelian quotients of the given group G . Unfortunately this approach is not sufficient. E.g. for the group $G = (C_3 \wr C_3) \times C_2$ we get $a(G) = 1/4, b(\mathbb{Q}, G) = 1, b(\mathbb{Q}(\zeta_3), G) = 2$. We can prove that there exist constants $c_1(G), c_2(G) > 0$ such that

$$c_1(G) x^{1/4} \leq Z(\mathbb{Q}, G; x) \leq c_2(G) x^{1/4} \text{ for } x \text{ large enough.}$$

Therefore this group does not contradict Conjecture 1.3. Similiar to our original example it is possible to choose $K = \mathbb{Q}(\zeta_3)$ as an intermediate extension, but this time it does not change the log-factor.

REFERENCES

- [1] Karim Belabas. Paramétrisation de structures algébriques et densité de discriminants [d'après Bhargava]. *Séminaire Bourbaki*, 56ème année(935), 2004.

- [2] Henri Cohen, Francisco Diaz y Diaz, and Michel Olivier. A survey of discriminant counting. In *Algorithmic number theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 80–94. Springer, Berlin, 2002.
- [3] Harold Davenport and Hans Arnold Heilbronn. On the density of discriminants of cubic fields. II. *Proc. Roy. Soc. London Ser. A*, 322(1551):405–420, 1971.
- [4] Jordan Ellenberg and Akshay Venkatesh. The number of extensions of a number field with fixed degree and bounded discriminant. arXiv:math.NT/0309153, 2003.
- [5] Jordan Ellenberg and Akshay Venkatesh. Counting extensions of function fields with bounded discriminant and specified Galois group. In *Geometric Methods in Algebra and Number Theory*, volume 235 in *Progress in Mathematics*, pages 151–168. Birkhäuser, 2005.
- [6] Jürgen Klüners and Gunter Malle. Counting nilpotent Galois extensions. *J. Reine Angew. Math.*, 572:1–26, 2004.
- [7] Gunter Malle. On the distribution of Galois groups. *J.Numb.Theory*, 92:315–322, 2002.
- [8] Gunter Malle. On the distribution of Galois groups II. *Exp.Math.*, 13:129–135, 2004.
- [9] David Wright. Distribution of discriminants of abelian extensions. *Proc. London Math. Soc.*, 58:17–50, 1989.

E-mail address: `klueners@mathematik.uni-kassel.de`

UNIVERSITÄT KASSEL, FACHBEREICH MATHEMATIK/INFORMATIK, HEINRICH-PLETT-STR. 40,
34132 KASSEL, GERMANY.